

Introduction to Vaccination, Presentation #1 Polio

[Introduction](#) [Sanitation](#)
[90% of Contageous Disease Mortality](#)
[Disappeared Before Vaccination](#)
[More mortality graphs](#) [Smallpox](#) [Polio](#) [Safe?](#)

1. Before the modern era, polio was a mild disease seldom resulting in paralysis.
2. The largest epidemic occurred about 1950 and had significantly declined by 1955 when vaccine was first introduced.
3. Disease can not be 'conquered' without addressing the disease cause.
4. Disease is not caused by a lack of "cow-pus", rotting material or carcinogenic poisons in the bloodstream.
5. Polio causes included:
 - pre-polio vaccines
 - nutritional imbalances
 - environmental poisons most notably pesticides

The fact that polio incidence increased after WWII and was only a minor problem before 1900 are clues that polio was a man made disease mainly caused by chemical pollution, most notably persistent pesticides which were used very carelessly. Sometimes parks were sprayed with DDT at the same time children were playing. Flies on milk cows were sprayed with not a thought of contaminating the cow's milk and thus poisoning children as a result.

[<Previous](#) [Next>](#)

6. Creation of and introduction of the Salk vaccine was motivated by fear that, without the vaccine, contributions to 'polio research' charities would drop.

Test results on the Salk vaccine were biased by throwing out polio cases 'after the first vaccination'. Test results were also biased by including only data from the '11 most accurate states' while ignoring the data from the other 33 states included in the test. What would the '33 least accurate states' data have shown?

7. Introduction of the Salk vaccine was beset first with known deaths related to the vaccine and then by a rise in polio cases.

8. Truth of the vaccine's true detriment was withheld from the public. Several states had rises in polio rate following vaccine introduction.

9. In the 3 years following the introduction of the vaccine:

The definition of epidemic was changed from 20 cases to 35 cases [per 100,000]!

The definition of polio was changed from paralysis occurring for 24 hours or more to paralysis occurring for 2 months!

The remaining polio cases were then diagnosed as meningitis!

[<Previous](#) [Next>](#)

**Hiding Polio:
After vaccination 'polio' became 'meningitis'**

Year	Polio	Meningitis	Total
July 1955	273	50	323
July 1961	65	161	226
July 1963	31	151	182
Sept. 1966	5	256	261

The above often quoted figures are credited to the Los Angeles County [USA] Health Index: Morbidity and Mortality, Reportable Diseases See:


<http://www.vaclib.org/basic/polio.htm>

For more hiding polio, see next graph:

[<Previous](#) [Next>](#)


Eradication of Poliomyelitis:
Progress in the Americas
Pediatr. Infect Dis. J. 1991;10:222-9
Vol 10, No. 3

Eradication of Poliomyelitis: As it was
presented



Note: The increasing sized black bars indicate larger numbers of 'reported' (Notified) cases of polio following vaccination. The decreasing sized grey bars indicate the fewer cases of polio were 'confirmed' thus allowing the false claim that the vaccine was working.

[<Previous](#) [Next>](#)



Source: Polio in the United States, Graphic Timeline: U.S. 1870-1998,
http://www.geocities.com/harpub/pol_all.htm


With 3 wonderful graphs relating DDT/etc to polio cases.

[<Previous](#) [Next>](#)

After thoughts on Polio:

In an effort to prevent a disease that caused 1000 deaths in the year before the vaccine introduction, SV40 contaminated vaccines are believed to now cause up to half of the 55,000 non-Hodgkin's lymphoma cases per year. [in the USA]

The following graph gives one an idea of what the world wide damage per year might be from the contaminated Polio vaccines.



source: *The Polio Hoax*

<http://christianparty.net/polio.htm>

<Previous Next>

**More lessons our schools forgot to tell us:
Diet Prevents Polio by Dr Sandler
(from: Whale.To Site)**

http://www.mercola.com/article/Diet/sugar/polio_sugar.htm

Sugar Increases Polio Risk -- Lessons For Other Viral Infections

A chapter from the book *Diet Prevents Polio* written by Benjamin P. Sandler, M.D., and published in 1951, at the height of the polio epidemic. In 1938, the only laboratory animal that could contract polio by experimental inoculation was the monkey. [Rabbits were completely immune to polio...] The next step was to lower the blood sugar of the rabbit to subnormal values with insulin injections, and then inoculate the rabbit with polio virus. This was done and it was found that the rabbits became infected and developed the disease.

See more at: <http://www.vaclib.org/basic/polio.htm>


**Another suppressed lesson?
Polio Cure**

The Origin of the 42-Year Stonewall of Vitamin C Landwehr, Journal of Orthomolecular Medicine, 1991 "Dr. F. R. Klenner, Reidsville, N.C.: It might be interesting to learn how poliomyelitis was treated in Reidsville, N.C., during the 1948 epidemic. In the past seven years, virus infections have been treated and cured in a period of seventy-two hours by the employment of massive frequent injections of ascorbic acid, or vitamin C."

See: www.seanet.com/~alexs/ascorbate/199x/index.htm#Landwehr-1991

[<Previous](#) [Next>](#)

The Complete Polio Story on One Page



Source: Polio in the United States, Graphic Timeline: U.S. 1870-1998,
http://www.geocities.com/harpub/pol_all.htm

With 3 wonderful graphs relating DDT/etc to polio cases.

[<Previous Safe?](#)